

Ticosonde CFH at Costa Rica: A seasonal climatology of tropical UT/LS water vapor and inter-comparisons with MLS and CALIPSO

Henry B. Selkirk, GESTAR/NASA GSFC

with

Holger Vömel, Deutscher Wetterdienst

Melody Avery, NASA LaRC

Karen Rosenlof, NOAA ESRL

Sean Davis and Dale Hurst, CIRES/NOAA ESRL

Mark Schoeberl, STC

Jorge Andrés Díaz, Univ. Costa Rica

Gary Morris, St. Edwards University



Aura STM 2014

Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder**
- B. Intercomparison to MLS v3.3 WV**
- C. Relationship between sonde saturation and CALIOP cloud fraction**
- D. Changes of WV in the tropics UT/LS**
 - Comparisons of sondes to MLS at 82 hPa
 - 2-km WV time series at Costa Rica



Aura STM 2014

Science drivers

- A. Processes controlling stratospheric water vapor: local freeze-drying, transport from remote tropical regions, in-mixing from higher latitudes**
- B. Continuing differences among *in situ* water vapor instruments**
- C. *Jensen et al.* [2013] found differing saturation regimes for low- and hi-particle density cirrus**
- D. Radiative forcing of UT/LS water vapor and impact in warming climate**



Aura STM 2014


Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder**
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS
 - Comparisons of sondes to MLS at 82 hPa
 - 2-km WV time series at Costa Rica




2005-2013

Frostpoint temperature, ambient temperature, and ozone



- ➡ Ozone gradient change nearly coincident with mean coldpoint
- ➡ Prominent inflection in O_3 profile @ 20 km




- Ozone gradient change well below mean coldpoint
- Smooth ozone profile throughout



Aura STM 2014

Tape recorder - I

Second half of year – JJA and SON




- 4 ppmv “hygropause” clearly visible at 468 K in JJA
- 5.5 ppmv maximum rises out of TTL in SON to 415 K



Aura STM 2014

Tape recorder - II

First half of year – DJF and MAM



- Profile minimum near trop in DJF and slow movement upward through MAM
- But SON max near 420 K has moved up to 460 K by DJF and 525 K in MAM
- Weak ascent in LMS during DJF and MAM, but more rapid above 450 K



Aura STM 2014

Talk topics


- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV**
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS
 - Comparisons of sondes to MLS at 82 hPa
 - 2-km WV time series at Costa Rica



Aura STM 2014

Coincidences with MLS v3.3


± 3 hours, $\Delta \leq 600$ km



December – February

65 coincident soundings:

Costa Rica (35), Biak (19), Kototabang (4), Tarawa (4)



June – September

28 coincident soundings:

Costa Rica (22), San Cristobál (6)



Aura STM 2014




Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction**
- D. Changes of WV in the tropics UT/LS
 - Comparisons of sondes to MLS at 82 hPa
 - 2-km WV time series at Costa Rica



Aura STM 2014


RH_{ice} : DJF vs. JJA






Aura STM 2014

CALIOP Vertical Feature Mask comparison 15 July 2005



VFM cloud analyzed at San José 10.5 to 14.5 km





CFH saturation layer 11.4 to 15 km

- 28 VFM/CFH coincidences closer than 400 km, 2006-2011
- Preliminary finding is that CALIOP is capturing most of the layers > 1 km deep
- However, VFM is just first step – ultimate goal is to compare to ice water content







Aura STM 2014

CALIOP Mean Cloud Fraction at San Jose, Costa Rica



Peak frequencies of sonde saturation






Aura STM 2014

Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS**
 - Comparisons of sondes to MLS at 82 hPa
 - 2-km WV time series at Costa Rica



Lower stratospheric water vapor MLS, frostpoint sondes and tropical CPT



Adapted from Fig. 2.43, BAMS 2014: State of the Climate 2013



Aura STM 2014


Time series calculations

- **Costa Rica CFH observations, July 2005 – July 2013**
- **Monthly time series of 2-km thick layers**
 - Linear average of 40 values at 50-m grid points
 - Centered at 3, 5, 7..... km
 - Layer average points computed only if 30 or more grid levels present
- **Multiple soundings in a month composited to a single value**
- **Monthly time means at each 1-km level derived for 8-year period**
 - Minimum sample size of 4 for each monthly level time mean
 - Exclude levels with 6 or more monthly time means missing
 - However, did allow interpolation of up to 2 consecutive missing months
 - Highest level meeting this criterion was 21 km
- **De-seasonalized anomaly time series generated**
 - Subtract long-term means for the month from each value at all levels meeting above criteria
- **Did linear fits in IGOR Pro to anomaly time series to check for trends**



Aura STM 2014


Layer-average WV time series and de-seasonalized anomalies





Aura STM 2014

Layer-average WV time series and de-seasonalized anomalies





Aura STM 2014

Summary

- Tape recorder structure at Costa Rica consistent with zonal mean, including slow uplift in early part of year in lowermost stratosphere
- Intercomparing MLS and CFH:
 - No real change from *Vömel et al.* [2007]: MLS drier than CFH in UT, wetter in LMS
 - Caveats: UT result swamped by large scatter; difference in LMS within instrumental uncertainties
- Seasonal mean CALIOP cloud fractions consistent with pattern of saturated sondes
 - Preliminary comparisons with coincident CALIOP feature mask data indicate that it can see most saturated sonde layers deeper than about 1 km are
 - However, addressing low- vs. high-density cirrus question will require information on ice water content
- Changes of WV in the tropics UT/LS:
 - Ticosonde CFH following tropical CPT values
 - Weak upward trends over nine years of program not significant – which is not unexpected given dominance of interannual variability